


ISSN (E): 2708-2601

ISSN (P): 2708-2598

Medical Journal of South Punjab

Article DOI:10.61581/MJSP.VOL06/02/05

Volume 6, Issue 2, 2025



Increase in Vessel Luminal Diameter after Rotational Atherectomy: A study from Multan Institute of Cardiology, Pakistan

Publication History

Received: Jan 27, 2025 Revised: Mar 23, 2025

Accepted: May 01, 2025 Published: June 30, 2025

Authors and Affiliation:

Momin Rasheed¹, Badar Ul Ahad Gill², Muhammad Ikram Farid³, Mubashir Sherwani⁴, Muhammad Shahzad Fareed⁵, Masood Ahmad Khan⁶

¹⁻⁶Chaudhry Pervaiz Elahi Institute of Cardiology, Multan, Pakistan.

*Corresponding Author Email:

mominrasheed@hotmail.com

Copyright & Licensing:



Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a [Creative Commons Attribution \(CC-BY\) 4.0 License](https://creativecommons.org/licenses/by/4.0/) that allows others to share the work with an acknowledgment of the work's authorship and initial publication in this journal.

Conflict of Interest:

Author(s) declared no conflict of interest.

Acknowledgment:

No Funding received.

Citation: Rasheed M, Gill BUA, Farid MI, Sherwani M, Fareed MS, Khan MA. Increase in Vessel Luminal Diameter after Rotational Atherectomy: A study from Multan Institute of Cardiology, Pakistan. Medical Journal of South Punjab. 2025 June 30; 6(2):28-34.

Please scan me to access online.



An official publication of

Medteach Private Limited, Multan, Pakistan.

Email: farman@mjsp.com.pk, Website: <https://mjsp.com.pk/index.php/mjsp>



Increase in Vessel Luminal Diameter after Rotational Atherectomy: A study from Multan Institute of Cardiology, Pakistan

Momin Rasheed¹, Badar Ul Ahad Gill², Muhammad Ikram Farid³, Mubashir Sherwani⁴, Muhammad Shahzad Fareed⁵, Masood Ahmad Khan⁶

¹⁻⁶Chaudhry Pervaiz Elahi Institute of Cardiology, Multan, Pakistan.

*Corresponding Author Email: mominrasheed@hotmail.com

ABSTRACT

Objective: To determine the increase in vessel luminal diameter in patients undergoing rotational atherectomy at Multan Institute of Cardiology, Pakistan.

Methods: Our study enrolled 100 patients undergoing rotational atherectomy. After recording baseline characteristics, the pre- and post-procedural minimal luminal diameters and reference vessel diameters were noted. The change in luminal diameter was recorded in terms of mean and standard deviation. We stratified the data, applied a post-stratification t-test, and considered a *p*-value of less than 0.05 to be statistically significant.

Results: The mean age was 52.20 ± 11.19 years, with a higher proportion of male patients (76.0%). Risk factors included hypertension (45.0%), smoking (40.0%), diabetes (34.0%), and hyperlipidemia (32.0%). The most frequently involved lesion was the left anterior descending artery (LAD: 55.0%), with mid-distal location (65.0%) and severe calcification (71.0%). The minimal lumen diameter (MLD) significantly increased from pre-procedure and post-procedure 0.75 ± 0.14 mm to 1.88 ± 0.22 mm ($p < 0.001$). At the same time, the mean luminal gain was 1.13 ± 0.026 mm. The luminal gain with relation to reference diameter was 46%.

Conclusion: Rotational atherectomy can lead to significant luminal gain with remarkable procedural success.

Keywords : Rotablation; Rotational Atherectomy; Rotational Angioplasty; Luminal Diameter; Vessel Diameter

1. INTRODUCTION

Despite considerable progress in percutaneous coronary intervention (PCI) over the last four decades, severe calcification within atherosclerotic plaque remains the principal factor contributing to adverse clinical outcomes in patients.^{1,2} The incidence of significant coronary calcification is increasing, linked to an aging demographic.^{3,4} Significant calcifications in coronary arteries pose a major challenge to the successful implantation and expansion of stents. Intracoronary imaging is crucial for the identification, localization, and quantification of coronary calcification.⁵ Planning for PCI is recommended, especially in cases where substantial calcification is expected.⁶ Interventional devices intended to alter calcified lesions before balloon angioplasty and stenting are classified into two techniques: atherectomy and non-atherectomy techniques. Lesion modification options that exclude atherectomy include balloon-based procedures, such as cutting/scoring balloons, semi-compliant balloons, ultra-high-pressure balloons, and intravascular lithotripsy.

Rotational atherectomy (RA), or rotablation, was introduced three decades ago as a treatment for complex CAD characterised by severely calcified lesions.⁷ In 2015, the UK had the incidence of RA procedure during PCI (3.1%), despite >20% of highly calcified coronary lesions being treated with PCI throughout Europe^{5,8}. International rates range from 0.8% to 3.1%. The absence of standardised procedures, inadequately structured training, and challenges associated with procedural complexity impede the broader adoption of RA^{9,10}.

In a study in Pakistan, calcification was mild to moderate in 16.4% and severe in 9.6% of patients, indicating a notable presence of vascular calcification in this group¹¹. The prevalence of patients who require rotablation is significantly high. However, the data regarding the rate of

estimated procedures undergoing rotablation is lacking. Outcomes and characteristics of RA in our patients are expected to differ due to increased age, frailty, multiple co-morbidities, and complex lesions¹². Additionally, numerous studies conducted on Western populations have demonstrated the effectiveness of RA in improving vascular patency and procedural success^{13,14}. Nevertheless, data concerning South Asian populations, particularly from Pakistan, are scarce, despite the rising incidence of complex coronary lesions in this area¹⁵. This study was conducted at the Multan Institute of Cardiology, Pakistan, to evaluate the increase in vessel luminal diameter in patients undergoing RA. This evaluation is essential for understanding procedural outcomes and guiding optimal interventional strategies tailored to the local patient population.

2. METHODOLOGY

Following institutional ethical approval (ERB# 122, dated April 9, 2024), a descriptive study was conducted at the Cardiology Department of CPEIC, Multan, from April 10, 2024, to April 10, 2025. One hundred patients were enrolled, as the minimum sample size required was 85, calculated using the WHO calculator, with an absolute precision of 10% and a 33% increase in luminal diameter. Patients 18 years and older, undergoing elective PCI with rotational atherectomy for severely calcified coronary lesions, having de novo coronary artery disease (CAD) with significant stenosis (>70%) in at least one major epicardial vessel, lesions considered undilatable or high resistance to balloon angioplasty, requiring plaque modification, reference vessel diameter between 2.5 mm and 4.0 mm, and those who are unwilling or not ideal candidates for CABG were included in the study. Patients with severe left ventricular dysfunction (LVEF <30%), unprotected left primary disease requiring atherectomy, severe renal impairment (eGFR <30 mL/min/1.73 m²) or

on dialysis, active bleeding or contraindications to antiplatelet therapy, severe allergy to contrast media or medications used during PCI, Pregnancy or lactation, and in-stent restenosis (ISR) requiring repeat RA were excluded from the study.

After obtaining informed written consent, the study included a minimum of 100 patients who fulfilled the inclusion criteria. Baseline characteristics, including age, gender, DM, HTN, smoking, hyperlipidemia, culprit vessel, lesion location, and lesion calcification severity, were noted. The minimal luminal diameter (MLD) and reference vessel diameter were measured before the procedure using quantitative coronary angiography (QCA) or intravascular imaging. After coronary catheter engagement and lesion wiring according to standard protocol, a Rotablator system (Boston Scientific) was used during the RA procedure. The burr size was selected based on the lesion and vessel size, usually ranging between 1.25 mm and 1.75 mm. Rotational speed was maintained between 140,000 and 180,000 rpm. Multiple short burr runs were performed, each lasting no more than 20 seconds, with adequate flushing between passes to minimise complications. Adjunctive therapies, such as cutting balloons or non-compliant balloon angioplasty, were used when necessary to optimize lesion preparation. Stenting was performed according to the standard protocol, and post-dilatation was documented. Procedural parameters such as fluoroscopy time and contrast volume were documented. Post-procedure, the final MLD was measured to assess luminal gain. Angiographic success was defined as achieving TIMI grade 3 flow with residual stenosis <30%. Complications such as coronary dissection, perforation, slow flow/no-reflow, bleeding, or periprocedural myocardial infarction were monitored and recorded.

Data was analysed using SPSS v.26. Qualitative variables, such as gender, DM, HTN, smoker, hyperlipidemia, culprit vessel, lesion location (proximal/mid/distal),

lesion calcification (diffuse/non-diffuse), and procedural success, were measured in frequency and percentages. Quantitative variables, including age, minimal luminal diameter (pre- and post-procedure), minimum reference vessel diameter (pre- and post-procedure), and TIMI flow grade (pre- and post-procedure), were measured in terms of mean and standard deviation. The luminal gain was recorded in terms of mean, standard deviation, and percentage, using the formula: $[(\text{post-MLD} - \text{Pre-MLD}) / \text{mean pre RVD}]$. Pre- and post-MLD, RVD were compared, and a p-value of ≤ 0.05 was considered statistically significant.

3. RESULTS

The study included 100 patients who met the inclusion criteria. The mean age was 52.20 ± 11.19 years. There were more male (76.0%) patients. Baseline comorbidities included hypertension (45.0%), smoking (40.0%), diabetes (34.0%), and hyperlipidemia (32.0%). Lesions most frequently involved the left anterior descending artery (LAD: 55.0%), followed by the right coronary artery (RCA: 29.0%), with mid-distal location (65.0%) and severe calcification (71.0%). Procedural parameters included a mean lesion length of 22.29 ± 5.22 mm, burr size of 1.58 ± 0.22 mm, and rotational speed of $178,323 \pm 6,867.63$ RPM. (Table. I). The mean proximal reference diameter decreased from pre-procedure and post-procedure 2.58 ± 0.35 mm to 2.39 ± 0.11 mm ($p < 0.001$). The minimal lumen diameter (MLD) significantly increased from pre-procedure and post-procedure 0.75 ± 0.14 mm to 1.88 ± 0.22 mm ($p < 0.001$). The distal reference diameter decreased from pre-procedure and post-procedure 2.30 ± 0.32 mm to 2.01 ± 0.56 mm ($p < 0.001$). (Table. II). Procedural success was observed as (96.0%). (Figure. I). At the same time, the mean luminal gain was 1.13 ± 0.026 mm. The luminal gain with relation to reference diameter was 46%.


Table-I:
Demographics, baseline, and lesion characteristics of the patients

Characteristics	N (%)	Mean ± S.D
Demographics characteristics		
Age (years)		52.20±11.19
≤40	23 (23.0)	
>40	77 (77.0)	
Gender		
Male	76 (76.0)	
Female	24 (24.0)	
Baseline characteristics		
Diabetes	34 (34.0)	
Hypertension	45 (45.0)	
Smoking	40 (40.0)	
Hyperlipidemia	32 (32.0)	
Lesion characteristics		
Culprit vessel		
LAD	55 (55.0)	
RCA	29 (29.0)	
LCX	9 (9.0)	
LM to LAD/LCX	7 (7.0)	
Lesion location		
Mid-distal	65 (65.0)	
Osteo-proximal	35 (35.0)	
Calcification		
Mild to moderate	29 (29.0)	
Severe	71 (71.0)	
Lesion length (mm)		22.29±5.22
Burr Size (mm)		1.58±0.22
Rotation Per Minute		178323±6867.63

Table-II:
Comparison of Pre-procedure and post-procedure vessel diameters

Variable	Pre-procedure (Mean \pm S.D)	Post-procedure (Mean \pm S.D)	Mean difference	t-test value	p-value
Proximal reference (mm)	2.58 \pm 0.35	2.39 \pm 0.11	-0.19	5.32	<0.001
MLD (mm)	0.75 \pm 0.14	1.88 \pm 0.22	1.13	-40.79	<0.001
Distal reference (mm)	2.30 \pm 0.32	2.01 \pm 0.56	-0.29	4.11	<0.001

Figure-I
Procedural success of the patients



4. DISCUSSION

Atherectomy techniques, like rotational and orbital methods, utilize a rapidly spinning burr to modify calcified plaques. RA is performed in various conditions, encompassing high-risk patients and complex anatomical structures. The device utilizes a diamond-encrusted elliptical burr to ablate calcified plaque, rotating at speeds ranging from 140,000 to 180,000 rpm via a helical drive shaft. This has progressed to a calcified lesion, utilizing a specialized guidewire. A single burr passage generally suffices to refine the arterial lumen and eliminate intravascular calcium rings, thus aiding the passage and dilation of balloons for stent implantation.

In our study, the mean age of the population undergoing rota ablation was 52.20 \pm 11.19 years. Most patients were male (76.0%). Risk factors included hypertension (45.0%), smoking (40.0%), diabetes (34.0%), and hyperlipidemia (32.0%). In most studies, the mean age of the population is around 69-75 years,^{16,17} which is older than ours, indicating changes in calcified lesions at an early age, likely due to an increase in diabetes.¹⁸ A separate study indicated that systolic blood pressure (BP) \geq 140 mmHg, extended lesion length, acute angulation, elevated initial burr-to-artery ratio, and the use of beta blockers were adverse prognostic indicators in patients undergoing RA¹⁹.

In our study, the lesions most commonly undergoing RA were LAD (55%), with mid-distal location (65.0%) and severe calcification (71.0%). Procedural parameters included a mean lesion length of 22.29 \pm 5.22 mm, burr size of 1.58 \pm 0.22 mm, and rotational speed of 178,323 \pm 6,867.63 RPM. These findings were consistent with a study by Chen et al., in which the LAD was involved in 57% of patients. The mid-distal location was observed in 60% of cases, and heavy calcification was present in 80% of the population. In $>50\%$ patients, burr size of 1.5

used²⁰. Another study reported 73% of heavy calcifications in RA patients²¹. Procedural success was observed as (96.0%). At the same time, the mean luminal gain was 1.13 ± 0.026 mm (46%). The procedural success rate of the CTO-vessel undergoing RA was 93%, as reported by Ayoub et al.²² The procedural success was 97% in a study by Khan et al.¹². The luminal gain in ISR patients undergoing RA was 41.3% in a survey conducted by Sayegh et al.²³. At the same time, the luminal gain was 1.34 mm, as reported by Farhat et al.²⁴ Our study has a few limitations, including being a single-center study with a relatively small sample size. Moreover, it was just a descriptive study with no MACES on follow-up. Further studies with a like comparison of Rota-ablation efficacy at different speeds, burr sizes, and measurement of vessel size using advanced modalities like IVUS are recommended.

5. CONCLUSION

Our study concludes that rotational atherectomy can result in significant luminal gain of up to 46% and remarkable procedural success of up to 96%. This study demonstrates that RA significantly increases the vessel luminal diameter, facilitating optimal stent delivery and expansion. The findings from our study validated the procedural efficacy and safety of RA in the local population, supporting its use in complex percutaneous coronary interventions (PCIs).

6. REFERENCES

1. Shi X, Gao J, Lv Q, Cai H, Wang F, Ye R, et al. Calcification in atherosclerotic plaque vulnerability: Friend or foe? *Front Physiol.* 2020;11:56.
2. Sugane H, Kataoka Y, Otsuka F, Nakaoku Y, Nishimura K, Nakano H, et al. Cardiac outcomes in patients with acute coronary syndrome attributable to calcified nodule. *Atherosclerosis.* 2021;318:70–5.
3. Graby J, Soto-Hernaez J, Murphy D, Oldman JLAF, Burnett TA, Charters PFP, et al. Coronary artery calcification on routine CT has prognostic and treatment implications for all ages. *Clin Radiol.* 2023 Jun;78(6):412–420.
4. Yi B, Zeng W, Lv L, Hua P. Changing epidemiology of calcific aortic valve disease: 30-year trends of incidence, prevalence, and deaths across 204 countries and territories. *Aging (Albany NY).* 2021;13(9):12710–32.
5. Khattak S, Sharma H, Khan SQ. Atherectomy techniques: Rotablation, orbital and laser. *Interv Cardiol.* 2024;19:e21.
6. Barbato E, Gallinoro E, Abdel-Wahab M, Andreini D, Carrié D, Di Mario C, et al. Management strategies for heavily calcified coronary stenoses: an EAPCI clinical consensus statement in collaboration with the EURO4C-PCR group. *Eur Heart J.* 2023;44(41):4340–56.
7. Allali A, Abdel-Wahab M, Elbasha K, Mankerious N, Traboulsi H, Kastrati A, et al. Rotational atherectomy of calcified coronary lesions: current practice and insights from two randomized trials. *Clin Res Cardiol.* 2023;112(9):1143–63.
8. Sharma V, Abdul F, Haider ST, Din J, Talwar S, O'kane P, et al. Rotablation in the very elderly-safer than we think? *Cardiovascular Revascularization Medicine.* 2021;22:36–41.
9. Maier A, Gissler MC, Jäckel M, Oettinger V, Bacmeister L, Heidenreich A, et al. Procedural safety of rotational atherectomy and modified balloon angioplasty: insights from a German national registry.
10. Xu J, Wang YW, Ma LK, Hu H, Chen HW, Hua JS, et al. Safety and efficacy of different rotational speed during

- rotational atherectomy in coronary heart disease patients (RACE): study protocol for a randomized controlled trial. *Trials*. 2025;26(1):126.
11. Imtiaz S, Ahmed DS, Tariq DS, Chughtai DJ, Saeed DF, Memon S, et al. Vascular Calcification in maintenance Hemodialysis patients in Pakistan – A single center Cross-sectional study. *Pak J Kidney Dis*. 2022;6(4):16–25.
 12. Khan BK, Faheem O, Rahman MN. Characteristics and outcomes of rotational atherectomy in a tertiary care cardiology facility in Pakistan. *Pak Hear J*. 2021 [cited 2025 May 7];54(3):214–8.
 13. Tomasiewicz B, Kubler P, Zimoch W, Kosowski M, Wańha W, Ładziński S, et al. Acute angulation and sequential lesion increase the risk of rotational atherectomy failure. *Circ J*. 2021;85(6):867–76.
 14. Blachutzik F, Meier S, Weissner M, Schlattner S, Gori T, Ullrich-Daub H, et al. Comparison of coronary intravascular lithotripsy and rotational atherectomy in the modification of severely calcified stenoses. *Am J Cardiol*. 2023;197:93–100.
 15. Khan S, Shah SHA, Basir W. Patterns of coronary artery occlusion in acute coronary syndrome patients undergoing coronary angiography in a cardiac center of a low-middle income country. *Pak Hear J*. 2025;58(1):73–8.
 16. Hemetsberger R, Mankerious N, Muntané-Carol G, Temporal J, Sulimov D, Gaede L, et al. In-hospital outcomes of rotational atherectomy in ST-elevation myocardial infarction: Results from the multicentre ROTA-STEMI network. *Can J Cardiol*. 2024;40(7):1226–33.
 17. Mahorkar A, Mahorkar V, Mahorkar A, Mahorkar U, Namdeo T, Sarwale S, et al. Comparative Outcomes of Rota Ablation in Calcific Coronary Lesions: Impact of SYNTAX Score, LV Dysfunction and Imaging vs. Non-Imaging-Guided PCI-A Single-Centre Experience from Central India. *Journal of Cardiovascular Disease Research*. 2024;15(11):278–91.
 18. Khan MI, Naseem M, Khalil MSUD, Yar A, Khan MR, Orakzai MH. Correlation between duration of diabetes and severity of coronary artery disease in patients undergoing coronary angiography at a tertiary care hospital. *Pak Hear J*. 2024;57(4):329–36.
 19. Sakakura K, Taniguchi Y, Yamamoto K, Tsukui T, Jinnouchi H, Seguchi M, et al. Modifiable and unmodifiable factors associated with slow flow following rotational atherectomy. *PLoS One*. 2021;16(4):e0250757.
 20. Chen YW, Lai CH, Su CS, Chang WC, Wang CY, Chen WJ, et al. The procedural and clinical outcomes of rotational atherectomy in patients presenting with acute myocardial infarction. *Front Cardiovasc Med*. 2022;9:846564.
 21. Khan AA, Murtaza G, Khalid MF, White CJ, Mamas MA, Mukherjee D, et al. Outcomes of rotational atherectomy versus orbital atherectomy for the treatment of heavily calcified coronary stenosis: A systematic review and meta-analysis. *Catheter Cardiovasc Interv*. 2021;98(5):884–92.
 22. Ayoub M, Corpataux N, Behnes M, Schupp T, Forner J, Akin I, et al. Safety and efficiency of rotational atherectomy in chronic total coronary occlusion-one-year clinical outcomes of an observational registry. *J Clin Med*. 2023;12(10).
 23. Sayegh M, Proczka M, Mancuso D, Huntress L, Savarese J, Ricotta J, et al. Effects of atherectomy on immediate luminal gain in patients with in-stent restenosis (ISR) with or without Stent

- thrombosis. J Vasc Surg. 2024;79(6):e178.
- 24.** Farhat H, Kuzemczak M, Durel N, Caillot N, Pawłowski T, Lipiecki J. Rotational atherectomy versus intravascular lithotripsy for calcified in-Stent restenosis: A single-center study with 1-year follow-up. Am J Cardiol. 2023;205:413–9.